

## Inecuatii trigonometrice

Metoda principala de rezolvare a inecuatilor trigonometrice consta in reducerea lor la inecuatii de forma

$$\sin x \vee a, \quad \cos x \vee a, \quad \operatorname{tg} x \vee a, \quad \operatorname{ctg} x \vee a, \quad (1)$$


unde  $a \in \mathbf{R}$ , semnul " $\vee$ " desemneaza semnul compararii si inlocuieste oricare din semnele " $>$ ", " $\geq$ ", " $<$ ", " $\leq$ " si utilizarea afirmatiilor ce urmeaza.

**Afirmatia 1.** Multimea solutiilor inecuatiei

$$\sin x > a \quad (2)$$

este

1.  $\mathbf{R}$ , daca  $a < -1$ ;
2.  $\bigcup_{k \in \mathbf{Z}} (\arcsin a + 2\pi k; \pi - \arcsin a + 2\pi k)$ , daca  $-1 \leq a < 1$ ;
3. Multimea vida, daca  $a \geq 1$ .




**Afirmatia 2.** Multimea solutiilor inecuatiei

$$\sin x < a \quad (3)$$

este

1.  $\mathbf{R}$ , daca  $a > 1$ ;
2.  $\bigcup_{k \in \mathbf{Z}} (-\pi - \arcsin a + 2\pi k; \arcsin a + 2\pi k)$ , daca  $-1 < a \leq 1$ ;
3. Multimea vida, daca  $a \leq -1$ .




**Afirmatia 3.** Multimea solutiilor inecuatiei

$$\cos x > a \tag{4}$$

este

1.  $\mathbf{R}$ , daca  $a < -1$ ;
2.  $\bigcup_{k \in \mathbf{Z}} (2\pi k - \arccos a; 2\pi k + \arccos a)$ , daca  $-1 \leq a < 1$ ;
3. Multimea vida, daca  $a \geq 1$ .




**Afirmatia 4.** Multimea solutiilor inecuatiei

$$\cos x < a \tag{5}$$

este


1.  $\mathbf{R}$ , daca  $a > 1$ ;
2.  $\bigcup_{k \in \mathbf{Z}} (2\pi k + \arccos a; 2\pi(k + 1) - \arccos a)$ , daca  $-1 < a \leq 1$ ;
3. Multimea vida, daca  $a \leq -1$ .



**Afirmatia 5.** Multimea solutiilor inecuatiei

$$\operatorname{tg} x > a \quad (6)$$


este  $\bigcup_{k \in \mathbf{Z}} (\operatorname{arctg} a + \pi k; \frac{\pi}{2} + \pi k)$ .



**Afirmatia 6.** Multimea solutiilor inecuatiei

$$\operatorname{tg} x < a \quad (7)$$


este  $\bigcup_{k \in \mathbf{Z}} (-\frac{\pi}{2} + \pi k; \operatorname{arctg} a + \pi k)$ .



**Afirmatia 7.** Multimea solutiilor inecuatiei

$$\operatorname{ctg} x > a \quad (8)$$


este  $\bigcup_{k \in \mathbf{Z}} (\pi k; \operatorname{arctg} a + \pi k)$ .



**Afirmatia 8.** Multimea solutiilor inecuatiei

$$\operatorname{ctg} x < a \tag{9}$$

este  $\bigcup_{k \in \mathbf{Z}} (\operatorname{arctg} a + \pi k; \pi(k + 1))$



**Nota.** 1. Daca semnul inegalitatii in (2)-(9) nu este strict, in multimea solutiilor inecuatiiilor se includ si solutiile ecuatiei respective.

2. Afirmatiile 1-8 se obtin nemijlocit analizand graficul functiilor trigonometrice respective.

**Exemplul 1.** Sa se rezolve inecuatiile

- |  |   |
|--|---|
| 1) $\sin 2x < \frac{1}{2}$ ;                                 | 7) $\operatorname{ctg}^2 x - \operatorname{ctg} x - 2 \leq 0$ ;                                     |
| 2) $2 \sin \left( \frac{\pi}{4} - x \right) \leq \sqrt{2}$ ; | 8) $\sin 2x - \sqrt{3} \cos 2x > \sqrt{2}$ ;  |
| 3) $\cos^2 x \geq \frac{1}{4}$ ;                             | 9) $\frac{2 \operatorname{tg} x}{1 + \operatorname{tg} x} + \frac{1}{\operatorname{tg} x} \geq 2$ ; |
| 4) $-2 \leq \operatorname{tg} x < 1$ ;                       | 10) $4 \sin x \cos x (\cos^2 x - \sin^2 x) < \sin 6x$ ;   |
| 5) $2 \sin^2 x - 5 \sin x + 2 > 0$ ;                         | 11) $\sin x \sin 3x \geq \sin 5x \sin 7x$ ;   |
| 6) $\sin^4 x + \cos^4 x \geq \frac{\sqrt{3}}{2}$ ;           | 12) $\sin x + \sin 2x + \sin 3x > 0$ .  |

**Rezolvare.** 1) Se noteaza  $2x = t$  si se obtine inecuatia  $\sin t < \frac{1}{2}$  care, conform afirmatiei 2 are solutiile

$$2\pi k - \pi - \arcsin \frac{1}{2} < t < \arcsin \frac{1}{2} + 2\pi k, \quad k \in \mathbf{Z}.$$

Se revine la variabila initiala si, tinand seama ca  $\arcsin \frac{1}{2} = \frac{\pi}{6}$ , se obtine

$$2\pi k - \pi - \frac{\pi}{6} < 2x < \frac{\pi}{6} + 2\pi k, \quad k \in \mathbf{Z},$$

de unde

$$2\pi k - \frac{7\pi}{6} < 2x < \frac{\pi}{6} + 2\pi k, \quad k \in \mathbf{Z},$$

sau

$$\pi k - \frac{7\pi}{12} < x < \frac{\pi}{12} + \pi k, \quad k \in \mathbf{Z}.$$

Asadar, solutiile inecuatiei enuntate formeaza multimea

$$\bigcup_{k \in \mathbf{Z}} \left( \pi k - \frac{7\pi}{12}; \frac{\pi}{12} + \pi k \right).$$

2) Cum functia sinus este impară,

$$2 \sin \left( \frac{\pi}{4} - x \right) \leq \sqrt{2} \Leftrightarrow -2 \sin \left( x - \frac{\pi}{4} \right) \leq \sqrt{2} \Leftrightarrow \sin \left( x - \frac{\pi}{4} \right) \geq -\frac{1}{\sqrt{2}}.$$

Se noteaza  $t = x - \frac{\pi}{4}$  si se obtine inecuatia

$$\sin t \geq -\frac{1}{\sqrt{2}}$$

cu solutiile (a se vedea afirmatia 1 si nota 1)

$$2\pi k + \arcsin \left( -\frac{1}{\sqrt{2}} \right) \leq t \leq \pi - \arcsin \left( -\frac{1}{\sqrt{2}} \right) + 2\pi k, \quad k \in \mathbf{Z},$$

de unde, tinand seama ca  $\arcsin\left(-\frac{1}{\sqrt{2}}\right) = -\frac{\pi}{4}$ , se obtine

$$2\pi k - \frac{\pi}{4} \leq x - \frac{\pi}{4} \leq \pi + \frac{\pi}{4} + 2\pi k, \quad k \in \mathbf{Z},$$

sau

$$2\pi k \leq x \leq \frac{3\pi}{2} + 2\pi k, \quad k \in \mathbf{Z}.$$

3) Cum  $\cos^2 x = \frac{1 + \cos 2x}{2}$  inecuatiya devine  $\frac{1 + \cos 2x}{2} \geq \frac{1}{4}$  sau  $\cos 2x \geq -\frac{1}{2}$ . Se aplica afirmatia 3 si se obtine

$$2\pi k - \arccos\left(-\frac{1}{2}\right) \leq 2x \leq \arccos\left(-\frac{1}{2}\right) + 2\pi k.$$

Cum  $\arccos\left(-\frac{1}{2}\right) = \frac{2\pi}{3}$ , rezulta

$$2\pi k - \frac{2\pi}{3} \leq 2x \leq \frac{2\pi}{3} + 2\pi k, \quad k \in \mathbf{Z},$$

de unde

$$\pi k - \frac{\pi}{3} \leq x \leq \frac{\pi}{3} + \pi k, \quad k \in \mathbf{Z}.$$

Altfel,

$$\begin{aligned} \cos^2 x \geq \frac{1}{4} &\Leftrightarrow |\cos x| \geq \frac{1}{2} \Leftrightarrow \begin{cases} \cos x \geq \frac{1}{2}, \\ \cos x \leq -\frac{1}{2}, \end{cases} \Leftrightarrow \\ \Leftrightarrow \begin{cases} 2\pi n - \frac{\pi}{3} \leq x \leq \frac{\pi}{3} + 2\pi n, & n \in \mathbf{Z}, \\ 2\pi m + \frac{2\pi}{3} \leq x \leq \frac{4\pi}{3} + 2\pi m, & m \in \mathbf{Z} \end{cases} &\Leftrightarrow \pi k - \frac{\pi}{3} \leq x \leq \frac{\pi}{3} + \pi k, \quad k \in \mathbf{Z}. \end{aligned}$$

4) Se aplica afirmatiile 5 si 6 si se obtine

$$\begin{aligned} -2 \leq \operatorname{tg} x < 1 &\Leftrightarrow \begin{cases} \operatorname{tg} x < 1, \\ \operatorname{tg} x \geq -2, \end{cases} \Leftrightarrow \begin{cases} \pi n - \frac{\pi}{2} < x < \frac{\pi}{4} + \pi n, & n \in \mathbf{Z}, \\ \pi m - \operatorname{arctg} 2 < x < \frac{\pi}{2} + \pi m, & m \in \mathbf{Z}, \end{cases} \Leftrightarrow \\ &\Leftrightarrow \pi k - \operatorname{arctg} 2 \leq x < \frac{\pi}{4} + \pi k, \quad k \in \mathbf{Z}. \end{aligned}$$

5) Se noteaza  $t = \sin x$  si se obtine inecuatiya de gradul al doilea

$$2t^2 - 5t + 2 > 0$$

cu solutiile

$$\begin{cases} t < \frac{1}{2}, \\ t > 2, \end{cases}$$

de unde rezulta totalitatea de inecuatii

$$\begin{cases} \sin x > 2, \\ \sin x < \frac{1}{2}. \end{cases}$$

Prima inecuatie a totalitatii solutii nu are, iar din cea secunda se obtine

$$2\pi k - \frac{7\pi}{6} < x < \frac{\pi}{6} + 2\pi k, \quad k \in \mathbf{Z}.$$

6) Cum

$$\begin{aligned} \sin^4 x + \cos^4 x &= (\sin^2 x)^2 + (\cos^2 x)^2 = (\sin^2 x + \cos^2 x)^2 - 2\sin^2 x \cos^2 x = \\ &= 1 - \frac{1}{2} \sin^2 2x = 1 - \frac{1}{2} \cdot \frac{1 - \cos 4x}{2} = 1 - \frac{1 - \cos 4x}{4}, \end{aligned}$$

inecuatia devine

$$1 - \frac{1 - \cos 4x}{4} \geq \frac{\sqrt{3}}{2}$$

sau  $\cos 4x \geq 2\sqrt{3} - 3$ . Cum  $|2\sqrt{3} - 3| \leq 1$ , se aplica afirmatia 3 si se obtine

$$2\pi k - \arccos(2\sqrt{3} - 3) < 4x < \arccos(2\sqrt{3} - 3) + 2\pi k, \quad k \in \mathbf{Z},$$

sau

$$\frac{\pi k}{2} - \frac{1}{4} \arccos(2\sqrt{3} - 3) < x < \frac{1}{4} \arccos(2\sqrt{3} - 3) + \frac{\pi k}{2}, \quad k \in \mathbf{Z}.$$

7) Se noteaza  $t = \text{ctg } x$  si se obtine inecuatia patrata

$$t^2 - t - 2 \leq 0$$

cu solutiile  $-1 \leq t \leq 2$ , de unde  $-1 \leq \text{ctg } x \leq 2$ . Ultima inecuatie se rezolva utilizand afirmatiile 7 si 8:

$$-1 \leq \text{ctg } x \leq 2 \Leftrightarrow \begin{cases} \text{ctg } x \leq 2, \\ \text{ctg } x \geq -1, \end{cases} \Leftrightarrow \begin{cases} \pi k + \text{arcctg } 2 \leq x < \pi + \pi n, \quad n \in \mathbf{Z} \\ \pi m < x \leq \frac{3\pi}{4} + \pi m, \quad m \in \mathbf{Z} \end{cases} \Leftrightarrow$$

$$\Leftrightarrow \pi k + \text{arcctg } 2 \leq x \leq \frac{3\pi}{4} + \pi k, \quad k \in \mathbf{Z}.$$

8) Se utilizeaza metoda unghiului auxiliar si se obtine

$$\begin{aligned} \sin 2x - \sqrt{3} \cos 2x > \sqrt{2} &\Leftrightarrow \frac{1}{2} \sin 2x - \frac{\sqrt{3}}{2} \cos 2x > \frac{\sqrt{2}}{2} \Leftrightarrow \\ \Leftrightarrow \sin 2x \cos \frac{\pi}{3} - \cos 2x \sin \frac{\pi}{3} > \frac{\sqrt{2}}{2} &\Leftrightarrow \sin \left( 2x - \frac{\pi}{3} \right) > \frac{\sqrt{2}}{2} \Leftrightarrow \\ \Leftrightarrow 2\pi k + \frac{\pi}{4} < 2x - \frac{\pi}{3} < \pi - \frac{\pi}{4} + 2\pi k, \quad k \in \mathbf{Z} &\Leftrightarrow \\ \Leftrightarrow 2\pi k + \frac{\pi}{4} + \frac{\pi}{3} < 2x < \frac{3\pi}{4} + \frac{\pi}{3} + 2\pi k, \quad k \in \mathbf{Z} &\Leftrightarrow \\ \Leftrightarrow \pi k + \frac{7\pi}{24} < x < \frac{13\pi}{24} + \pi k, \quad k \in \mathbf{Z}. \end{aligned}$$

9) Se noteaza  $\operatorname{tg} x = t$  si se rezolva inecuatia in  $t$  utilizand metoda intervalelor:

$$\frac{2t}{1+t} + \frac{1}{t} \geq 2 \Leftrightarrow \frac{2t^2 + 1 + t - 2t(1+t)}{t(t+1)} \geq 0 \Leftrightarrow \frac{1-t}{t(t+1)} \geq 0 \Leftrightarrow \begin{cases} 0 < t \leq 1, \\ t < -1. \end{cases}$$

Asadar, se obtine totalitatea de inecuatii

$$\begin{cases} 0 < \operatorname{tg} x \leq 1, \\ \operatorname{tg} x < -1, \end{cases}$$


ce se rezolva, utilizand afirmatiile 5 si 6:

$$\begin{aligned} \begin{cases} 0 < \operatorname{tg} x \leq 1, \\ \operatorname{tg} x < -1, \end{cases} &\Leftrightarrow \begin{cases} \operatorname{tg} x \leq 1, \\ \operatorname{tg} x > 0, \end{cases} \Leftrightarrow \begin{cases} -\frac{\pi}{2} + \pi m < x < -\frac{\pi}{4} + \pi m, & m \in \mathbf{Z}, \end{cases} \\ &\Leftrightarrow \begin{cases} \pi n < x \leq \frac{\pi}{4} + \pi n, & n \in \mathbf{Z}, \\ -\frac{\pi}{2} + \pi m < x < -\frac{\pi}{4} + \pi m, & m \in \mathbf{Z}. \end{cases} \end{aligned}$$

10) Se utilizeaza formulele sinusului si cosinusului argumentului dublu si se obtine

$$\begin{aligned} 4 \sin x \cos x (\cos^2 x - \sin^2 x) < \sin 10x &\Leftrightarrow 2 \sin 2x \cdot \cos 2x < \sin 6x \Leftrightarrow \\ \Leftrightarrow \sin 4x < \sin 6x &\Leftrightarrow \sin 6x - \sin 4x > 0 \Leftrightarrow 2 \sin x \cos 5x > 0. \end{aligned}$$

Se tine seama ca  $2\pi$  este o perioada a functiei  $f(x) = \sin x \cos 5x$  si se utilizeaza metoda generalizata a intervalelor pentru un interval de lungime  $2\pi$ :



Astfel multimea solutiilor inecuatiei date este reuniunea multimilor

$$\begin{aligned} &\left(2\pi k; \frac{\pi}{10} + 2\pi k\right) \cup \left(\frac{3\pi}{10} + 2\pi k; \frac{\pi}{2} + 2\pi k\right) \cup \left(\frac{7\pi}{10} + 2\pi k; \frac{9\pi}{10} + 2\pi k\right) \cup \\ &\cup \left(2\pi k + \pi; \frac{11\pi}{10} + 2\pi k\right) \cup \left(2\pi k + \frac{13\pi}{10}; \frac{3\pi}{2} + 2\pi k\right) \cup \left(\frac{17\pi}{10} + 2\pi k; \frac{19\pi}{10} + 2\pi k\right). \end{aligned}$$

11)  $\sin x \sin 3x \geq \sin 2x \sin 4x \Leftrightarrow \frac{1}{2}(\cos 2x - \cos 4x) \geq \frac{1}{2}(\cos 2x - \cos 6x) \Leftrightarrow$   
 $\Leftrightarrow -\cos 4x \geq -\cos 6x \Leftrightarrow \cos 6x - \cos 4x \geq 0 \Leftrightarrow -2\sin x \sin 5x \geq 0 \Leftrightarrow \sin x \sin 5x \leq 0.$   
 Ultima inecuatie se rezolva similar exemplului precedent si se obtine multimea solutiilor

$$\bigcup_{k \in \mathbf{Z}} \left[ \frac{2\pi}{5}n; \frac{\pi}{5} + \frac{2\pi}{5}n \right].$$

12)  $\sin x + \sin 2x + \sin 3x > 0 \Leftrightarrow (\sin x + \sin 3x) + \sin 2x > 0 \Leftrightarrow 2\sin 2x \cos x + \sin 2x > 0 \Leftrightarrow$

$$\Leftrightarrow \sin 2x(2\cos x + 1) > 0 \Leftrightarrow \begin{cases} \sin 2x > 0, \\ \cos x > -\frac{1}{2}, \end{cases} \Leftrightarrow \begin{cases} \sin 2x < 0, \\ \cos x < -\frac{1}{2}, \end{cases}$$

$$\Leftrightarrow \begin{cases} \pi n < x < \frac{\pi}{2} + \pi n, & n \in \mathbf{Z}, \\ -\frac{2\pi}{3} + 2\pi m < x < \frac{2\pi}{3} + 2\pi m, & m \in \mathbf{Z}, \end{cases} \Leftrightarrow \begin{cases} \frac{\pi}{2} + \pi n < x < \pi + \pi n, & n \in \mathbf{Z}, \\ 2\pi m + \frac{2\pi}{3} < x < \frac{4\pi}{3} + 2\pi m, & m \in \mathbf{Z}, \end{cases}$$

$$\Leftrightarrow \begin{cases} 2\pi m < x < \frac{\pi}{2} + 2\pi m, & m \in \mathbf{Z}, \\ 2\pi m - \frac{2\pi}{3} < x < -\frac{\pi}{2} + 2\pi m, & m \in \mathbf{Z}, \\ 2\pi m + \frac{2\pi}{3} < x < \pi + 2\pi m, & m \in \mathbf{Z}. \end{cases}$$

### Exercitii pentru autoevaluare

Sa se rezolve inecuatiiile:

1.  $\operatorname{tg}^3 x + \operatorname{tg}^2 x - \operatorname{tg} x - 1 > 0;$
2.  $\operatorname{tg} x + \operatorname{ctg} x \leq 2;$
3.  $\sin 2x < \cos x;$
4.  $\cos x + \cos 2x + \cos 3x \geq 0;$
5.  $6\sin^2 x - 5\sin x + 1 > 0;$

6.  $\frac{2 \cos^2 x + \cos x - 1}{\sin x - 1} < 0;$
7.  $2 \cos\left(2x + \frac{\pi}{4}\right) - \sqrt{3} \leq 0;$
8.  $\operatorname{tg}\left(\frac{\pi}{4} - 2x\right) < -\sqrt{3};$
9.  $2 \sin^2 x + 9 \cos x - 6 \geq 0;$
10.  $\frac{\sin x}{1 + \cos x} \geq 0;$
11.  $4 \sin x \cos x - \sqrt{2} < 2(\sqrt{2} \cos x - \sin x);$
12.  $\cos 2x + \sin x \geq 0.$